

{rokbox title=|Epiactis georgiana :: Foto: Estefanía Rodríguez|
thumb=|images/stories/ieo/imagenespublicaciones/centro-oceanografico-baleares-covadonga-o
rejas-epiactis-2012-thumb.jpg|}images/stories/ieo/imagenespublicaciones/centro-oceanografico
-baleares-covadonga-orejas-epiactis-2012.jpg{/rokbox}

E. Rodríguez, **C. Orejas**, P. J. López-González, J. M. Gili, (2012). [Reproduction in the externally brooding sea anemone *Epiactis georgiana* in the Antarctic Peninsula and the Weddell Sea](#)
. Marine Biology. DOI 10.1007/s00227-012-2063-x.

Abstract: External parental care is uncommon among actinarians but common in *Epiactis* species. Here, several aspects of reproduction are analyzed for one of them,

Epiactis georgiana

. Samples were collected in December, January, February, March, and April in the Antarctic Peninsula and the eastern Weddell Sea, during 1998, 2000, 2002, and 2003. Most sexually mature individuals of

E. georgiana

are male or female, but some are hermaphrodites. This is the first report of hermaphroditism in

E. georgiana

, which is the third species of the genus with this sexual pattern. The results suggest that oogenesis starts in December and that at least two generations of oocytes overlap; a third generation is often brooded externally. Putative fertilization is likely internal, and larvae and/or embryos are externally brooded on the distal part of the adult column until an advanced developmental stage. Apparently *E. georgiana* reproduces seasonally, probably releasing the embryos/larvae in the last months of the austral spring (December). Inter-individual variability was observed in gametogenesis. In addition, specimens from the Antarctic Peninsula were larger than those from the Weddell Sea. This study represents the first step in understanding the reproductive mode of

E. georgiana

.

Keywords: Actinarians, *Epiactis*, reproduction, parental care, seasonal reproduction, Antarctica